

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/256805834>

An operator inequality and its consequences

ARTICLE in LINEAR ALGEBRA AND ITS APPLICATIONS · AUGUST 2013

Impact Factor: 0.94 · DOI: 10.1016/j.laa.2012.08.005

CITATIONS

3

READS

116

3 AUTHORS:



M. S. Moslehian

Ferdowsi University Of Mashhad

160 PUBLICATIONS 917 CITATIONS

SEE PROFILE



Jadranka Micic

University of Zagreb

30 PUBLICATIONS 118 CITATIONS

SEE PROFILE



Mohsen Kian

University of Bojnord

16 PUBLICATIONS 28 CITATIONS

SEE PROFILE

AN OPERATOR INEQUALITY AND ITS CONSEQUENCES

M. S. MOSLEHIAN, J. MIĆIĆ AND M. KIAN

Dedicated to Professor Harm Bart on the occasion of his 70th birthday

ABSTRACT. Let f be a continuous convex function on an interval J , let A, B, C, D be self-adjoint operators acting on a Hilbert space with spectra contained in J such that $A + D = B + C$ and $A \leq m \leq B, C \leq M \leq D$ for two real numbers $m < M$, and let Φ be a unital positive linear map on $\mathbb{B}(\mathcal{H})$. We prove the inequality

$$f(\Phi(B)) + f(\Phi(C)) \leq \Phi(f(A)) + \Phi(f(D)).$$

and apply it to obtain several inequalities such as the Jensen–Mercer operator inequality and the Petrović operator inequality.

1. INTRODUCTION AND PRELIMINARIES

Throughout the paper $\mathbb{B}(\mathcal{H})$ stands for the C^* -algebra of all bounded linear operators on a complex Hilbert space \mathcal{H} and I denotes the identity operator. The real subspace of $\mathbb{B}(\mathcal{H})$ consisting of all self-adjoint operators on \mathcal{H} is denoted by $\mathbb{B}(\mathcal{H})_h$. An operator A is said to be positive (denoted by $A \geq 0$) if $\langle Ax, x \rangle \geq 0$ for all $x \in \mathcal{H}$. If, in addition, A is invertible, then it is called strictly positive (denoted by $A > 0$). By $A \geq B$ we mean that $A - B$ is positive, while $A > B$ means that $A - B$ is strictly positive. A map Φ on $\mathbb{B}(\mathcal{H})$ is said to be positive if $\Phi(A) \geq 0$ for each $A \geq 0$ and is called unital if $\Phi(I) = I$.

Let f be a continuous real valued function defined on an interval J . The function f is called operator monotone if $A \geq B$ implies $f(A) \geq f(B)$ for all $A, B \in \mathbb{B}(\mathcal{H})_h$ with spectra in J . A function f is called operator convex on J if $f(\lambda A + (1 - \lambda)B) \leq \lambda f(A) + (1 - \lambda)f(B)$ or all $A, B \in \mathbb{B}(\mathcal{H})_h$ with spectra in J and all $\lambda \in [0, 1]$. If the function f is operator convex, then the so-called Jensen operator inequality $f(\Phi(A)) \leq \Phi(f(A))$ holds for any unital positive linear map Φ on $\mathbb{B}(\mathcal{H})$ and any $A \in \mathbb{B}(\mathcal{H})_h$ with spectrum contained in J . Many other

2010 *Mathematics Subject Classification.* 15A60, 47A63, 47A64, 26D15.

Key words and phrases. convex function, positive linear map, Jensen–Mercer operator inequality, Petrović operator inequality.

versions of the Jensen operator inequality can be found in [6]. Among them, the Jensen–Mercer operator inequality [8] reads as follows:

$$f\left(M + m - \sum_{i=1}^n \Phi_i(A_i)\right) \leq f(M) + f(m) - \sum_{i=1}^n \Phi_i(f(A_i)),$$

where f is a continuous convex function on an interval $[m, M]$, Φ_1, \dots, Φ_n are positive linear maps on $\mathbb{B}(\mathcal{H})$ with $\sum_{i=1}^n \Phi_i(I) = I$ and $A_1, \dots, A_n \in \mathbb{B}(\mathcal{H})_h$ with spectra contained in $[m, M]$.

Another interesting inequality is the Petrović inequality (see e.g. [10, page 152]). It states that if $f : [0, \infty) \rightarrow \mathbb{R}$ is a continuous convex function and A_1, \dots, A_n are positive operators such that $\sum_{i=1}^n A_i = MI$ for some scalar $M > 0$, then

$$\sum_{i=1}^n f(A_i) \leq f\left(\sum_{i=1}^n A_i\right) + (n-1)f(0).$$

If $f : [0, \infty) \rightarrow [0, \infty)$ is a convex function and $f(0) \leq 0$, then

$$f(a) + f(b) \leq f(a+b) \tag{1.1}$$

for all scalars $a, b \geq 0$. However, if the scalars a, b are replaced by two positive operators, this inequality may not hold. There have been many interesting works devoted to obtain norm or operator extensions of inequality (1.1) and other subadditivity inequalities. Ando and Zhan [1] showed that if f is a nonnegative operator monotone function on the interval $[0, \infty)$, then

$$|||f(A+B)||| \leq |||f(A) + f(B)||| \tag{1.2}$$

for all unitarily invariant norms $||| \cdot |||$ and all matrices $A, B \geq 0$. Bourin and Uchiyama [5] extended Ando-Zhan's result by showing that if $A, B \geq 0$ and f is a non-negative concave function on $[0, \infty)$, then (1.2) holds for all unitarily norms. Also Aujla and Bourin [3] showed that if $A, B \geq 0$ and $f : [0, \infty) \rightarrow [0, \infty)$ is a monotone concave function, then there exist unitaries U, V such that

$$f(A+B) \leq Uf(A)U^* + Vf(B)V^*.$$

The reader is referred to [4, 11, 7, 12, 9, 2] and references therein for recent information on the subject.

In their study of Hadamard's inequalities for co-ordinated convex functions, Hwang, Tseng and Yang proved that if $f : [a, b] \rightarrow \mathbb{R}$ is a convex function and

$a \leq y_1 \leq x_1 \leq x_2 \leq y_2 \leq b$ and $x_1 + x_2 = y_1 + y_2$, then

$$f(x_1) + f(x_2) \leq f(y_1) + f(y_2).$$

In this paper, we extend this inequality to operators acting on a Hilbert space and apply it to obtain a series of operator inequalities including the Jensen–Mercer operator inequality, the Petrović operator inequality.

2. RESULTS

We start this section with our main result.

Theorem 2.1. *Let f be a continuous convex function on an interval J . Let $A, B, C, D \in \mathbb{B}(\mathcal{H})_h$ with spectra contained in J such that $A + D = B + C$ and $A \leq m \leq B, C \leq M \leq D$ for two real numbers $m < M$. If Φ is a unital positive linear map on $\mathbb{B}(\mathcal{H})$, then*

$$f(\Phi(B)) + f(\Phi(C)) \leq \Phi(f(A)) + \Phi(f(D)). \quad (2.1)$$

If f is concave on J , then inequality (2.1) is reversed.

Proof. If $t \in [m, M]$, then $0 \leq \frac{M-t}{M-m} \leq 1$. Since f is convex on $[m, M]$, we have

$$f(t) = f\left(\frac{M-t}{M-m}m + \frac{t-m}{M-m}M\right) \leq \frac{M-t}{M-m}f(m) + \frac{t-m}{M-m}f(M). \quad (2.2)$$

If $t \in J \setminus (m, M)$, then $t \leq m$ or $M \leq t$. Suppose that $t \leq m$ (the case when $M \leq t$ can be verified similarly). Therefore $0 \leq \frac{M-m}{M-t} \leq 1$ and

$$f(m) = f\left(\frac{M-m}{M-t}t + \frac{m-t}{M-t}M\right) \leq \frac{M-m}{M-t}f(t) + \frac{m-t}{M-t}f(M),$$

whence

$$f(t) \geq \frac{M-t}{M-m}f(m) + \frac{t-m}{M-m}f(M) \quad (2.3)$$

for all $t \in J \setminus (m, M)$. Since $A \leq m$ and $D \geq M$, using functional calculus to inequality (2.3) we obtain

$$f(A) \geq \frac{M-A}{M-m}f(m) + \frac{A-m}{M-m}f(M) \quad (2.4)$$

and

$$f(D) \geq \frac{M-D}{M-m}f(m) + \frac{D-m}{M-m}f(M). \quad (2.5)$$

Applying the positive linear map Φ to both sides of (2.4) and (2.5) we get

$$\Phi(f(A)) \geq \frac{M-\Phi(A)}{M-m}f(m) + \frac{\Phi(A)-m}{M-m}f(M) \quad (2.6)$$

and

$$\Phi(f(D)) \geq \frac{M - \Phi(D)}{M - m} f(m) + \frac{\Phi(D) - m}{M - m} f(M). \quad (2.7)$$

Similarly, taking into account that $m \leq \Phi(B) \leq M$ and $m \leq \Phi(C) \leq M$ and using functional calculus to inequality (2.2) we obtain

$$f(\Phi(B)) \leq \frac{M - \Phi(B)}{M - m} f(m) + \frac{\Phi(B) - m}{M - m} f(M) \quad (2.8)$$

and

$$f(\Phi(C)) \leq \frac{M - \Phi(C)}{M - m} f(m) + \frac{\Phi(C) - m}{M - m} f(M). \quad (2.9)$$

Adding two inequalities (2.8) and (2.9) we obtain

$$\begin{aligned} f(\Phi(B)) + f(\Phi(C)) &\leq \frac{M - \Phi(B)}{M - m} f(m) + \frac{\Phi(B) - m}{M - m} f(M) \\ &\quad + \frac{M - \Phi(C)}{M - m} f(m) + \frac{\Phi(C) - m}{M - m} f(M) \\ &= \frac{2M - \Phi(B + C)}{M - m} f(m) + \frac{\Phi(B + C) - 2m}{M - m} f(M) \\ &= \frac{2M - \Phi(A + D)}{M - m} f(m) + \frac{\Phi(A + D) - 2m}{M - m} f(M) \\ &\quad \text{(by } A + D = B + C\text{)} \\ &= \frac{M - \Phi(A)}{M - m} f(m) + \frac{\Phi(A) - m}{M - m} f(M) \\ &\quad + \frac{M - \Phi(D)}{M - m} f(m) + \frac{\Phi(D) - m}{M - m} f(M) \\ &\leq \Phi(f(A)) + \Phi(f(D)) \quad \text{(by (2.6) and (2.7))} \end{aligned}$$

□

We give an example to clarify the situation in Theorem 2.1.

Example 2.2. Let the function f be defined on $[0, \infty)$ by $f(t) = t^3$ and the unital positive linear map $\Phi : \mathcal{M}_2(\mathbb{C}) \rightarrow \mathcal{M}_2(\mathbb{C})$ be defined by $\Phi(A) = (\frac{1}{2} \text{tr}(A))I$ for all Hermitian matrices $A \in \mathcal{M}_2(\mathbb{C})$. If

$$A = \begin{pmatrix} 1 & -1 \\ -1 & 1 \end{pmatrix}, \quad B = \begin{pmatrix} 4 & 1 \\ 1 & 4 \end{pmatrix}, \quad C = \begin{pmatrix} 6 & -1 \\ -1 & 7 \end{pmatrix}, \quad D = \begin{pmatrix} 9 & 1 \\ 1 & 10 \end{pmatrix},$$

then

$$0 \leq A < 3I \leq B \leq C \leq 8I < D \quad \text{and} \quad A + D = B + C,$$

whence

$$f(\Phi(B)) + f(\Phi(C)) = 338.625I \not\leq 897I = \Phi(f(A)) + \Phi(f(D)).$$

This shows that inequality (2.1) can be strict.

More generally, the next corollary gives other versions of inequality (2.1). The proof is similar to that of Theorem 2.1 and we omit it.

Corollary 2.3. *Let f be a continuous convex function on an interval J . Let $A_i, B_i, C_i, D_i \in \mathbb{B}(\mathcal{H})_h$ ($i = 1, \dots, n$) with spectra contained in J such that $A_i + D_i = B_i + C_i$ and $A_i \leq m \leq B_i \leq M \leq D_i$ and $A_i \leq m \leq C_i \leq M \leq D_i$ ($i = 1, \dots, n$). Let Φ_1, \dots, Φ_n be positive linear maps on $\mathbb{B}(\mathcal{H})$ with $\sum_{i=1}^n \Phi_i(I) = I$. Then*

$$\begin{aligned} (1) \quad & f\left(\sum_{i=1}^n \Phi_i(B_i)\right) + f\left(\sum_{i=1}^n \Phi_i(C_i)\right) \leq \sum_{i=1}^n \Phi_i(f(A_i)) + \sum_{i=1}^n \Phi_i(f(D_i)); \\ (2) \quad & \sum_{i=1}^n \Phi_i(f(B_i)) + \sum_{i=1}^n \Phi_i(f(C_i)) \leq f\left(\sum_{i=1}^n \Phi_i(A_i)\right) + f\left(\sum_{i=1}^n \Phi_i(D_i)\right); \\ (3) \quad & \sum_{i=1}^n \Phi_i(f(B_i)) + f\left(\sum_{i=1}^n \Phi_i(C_i)\right) \leq f\left(\sum_{i=1}^n \Phi_i(D_i)\right) + \sum_{i=1}^n \Phi_i(f(A_i)). \end{aligned}$$

The Jensen–Mercer operator inequality follows directly from Theorem 2.1.

Corollary 2.4. [8, Theorem 1] *Let Φ_1, \dots, Φ_n be positive linear maps on $\mathbb{B}(\mathcal{H})$ with $\sum_{i=1}^n \Phi_i(I) = I$ and $B_1, \dots, B_n \in \mathbb{B}(\mathcal{H})_h$ with spectra contained in $[m, M]$. If f is a continuous convex function on $[m, M]$, then*

$$f\left(m + M - \sum_{i=1}^n \Phi_i(B_i)\right) \leq f(m) + f(M) - \sum_{i=1}^n \Phi_i(f(B_i)).$$

Proof. Clearly $m \leq B_i \leq M$ ($i = 1, \dots, n$). Set $C_i = M + m - B_i$ ($i = 1, \dots, n$). Then $m \leq C_i \leq M$ and $B_i + C_i = m + M$ ($i = 1, \dots, n$). Applying inequality (3) of Corollary 2.3 when $A_i = mI$ and $D_i = MI$ we obtain

$$\sum_{i=1}^n \Phi_i(f(B_i)) + f\left(\sum_{i=1}^n \Phi_i(C_i)\right) \leq f(m) + f(M),$$

which is the desired inequality. □

The next result provides an extension of Petrović inequality.

Corollary 2.5. *Let $A, D, B_i \in \mathbb{B}(\mathcal{H})_h$ ($i = 1, \dots, n$) with spectra contained in an interval J such that $A + D = \sum_{i=1}^n B_i$ and $A \leq m \leq B_i \leq M \leq D$ ($i = 1, \dots, n$) for two real numbers $m < M$. If f is convex on J , then*

$$\sum_{i=1}^n f(B_i) \leq (n-1)f\left(\frac{1}{n-1}A\right) + f(D).$$

The next result is a Jensen operator inequality for continuous convex functions. First define the subset Ω of $\mathbb{B}_h(\mathcal{H}) \times \mathbb{B}_h(\mathcal{H})$ by

$$\Omega = \left\{ (A, B) \mid A \leq m \leq \frac{A+B}{2} \leq M \leq B, \text{ for some } m, M \in \mathbb{R} \right\}.$$

The authors would like to pose the following problem that is interesting on its own right.

Problem 2.6. Is there any characterization of Ω for either matrices or operators acting on infinite dimensional Hilbert spaces?

Corollary 2.7. *If f is a continuous convex function on an interval J , then*

$$f(\lambda A + (1-\lambda)D) \leq \lambda f(A) + (1-\lambda)f(D) \quad (2.10)$$

for all $(A, D) \in \Omega$ with spectra contained in J and all $\lambda \in [0, 1]$. If f is concave, then inequality (2.10) is reversed.

Proof. Letting Φ be the identity map and putting $C = B = \frac{A+D}{2}$ in inequality (2.1), we get

$$f\left(\frac{A+D}{2}\right) \leq \frac{f(A) + f(D)}{2}$$

for any $(A, D) \in \Omega$ with spectra contained in J which implies inequality (2.10). \square

Note that the existence of scalars $m < M$ is essential in Corollary 2.7, i.e., inequality (2.10) may not hold if $A, B \notin \Omega$.

Example 2.8. Consider the convex function $f(t) = t^3$ on $[0, \infty)$. Putting

$$A = \begin{pmatrix} 1 & 1 & 0 \\ 1 & 2 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad B = \begin{pmatrix} 1 & 1 & 0 \\ 1 & 4 & 0 \\ 0 & 0 & 2 \end{pmatrix}$$

we have $0 \leq A \leq B$. There is no scalar m such that $A \leq m \leq \frac{A+B}{2}$. Now

$$\left(\frac{A+B}{2}\right)^3 = \begin{pmatrix} 6 & 14 & 0 \\ 14 & 34 & 0 \\ 0 & 0 & 3.375 \end{pmatrix} \not\leq \begin{pmatrix} 6 & 15 & 0 \\ 15 & 43 & 0 \\ 0 & 0 & 4.5 \end{pmatrix} = \frac{A^3 + B^3}{2},$$

shows that not only f is not operator convex but also the existence of scalars m, M are essential in definition of Ω .

Generally, inequality (1.1) would be false if we replace scalars a, b with two positive operators. To see this consider the convex function $f(t) = t^3$ on $[0, \infty)$ and two positive matrices A, B in Example 2.8. The following corollary gives an operator version of (1.1). The reader may compare it with [7, Theorem 2.2].

Corollary 2.9. *If $f : [0, \infty) \rightarrow [0, \infty)$ is a continuous convex function with $f(0) \leq 0$, then*

$$f(A) + f(B) \leq f(A + B)$$

for all strictly positive operators A, B for which $A \leq M \leq A + B$ and $B \leq M \leq A + B$ for some scalar M .

Proof. It follows directly from Corollary 2.5. □

Theorem 2.10. *Let f be a continuous function on an interval J . Let $A, B, C, D \in \mathbb{B}(\mathcal{H})_h$ with spectra contained in J such that $A \leq m \leq B, C \leq M \leq D$ for two real numbers $m \leq M$. If f is convex and one of the following conditions*

- (i) $B + C \leq A + D$ and $f(m) \leq f(M)$
- (ii) $A + D \leq B + C$ and $f(M) \leq f(m)$

is satisfied, then

$$f(B) + f(C) \leq f(A) + f(D). \tag{2.11}$$

If f is concave and one of the following conditions

- (iii) $B + C \leq A + D$ and $f(M) \leq f(m)$
- (iv) $A + D \leq B + C$ and $f(m) \leq f(M)$

is satisfied, then inequality (2.11) is reversed.

Proof. Let f be convex and (i) is valid. It follows from (2.2) that

$$f(B) \leq \frac{f(M) - f(m)}{M - m}B + \frac{f(m)M - f(M)m}{M - m}$$

and

$$f(C) \leq \frac{f(M) - f(m)}{M - m}C + \frac{f(m)M - f(M)m}{M - m}.$$

Summing above inequalities we get

$$\begin{aligned} f(B) + f(C) &\leq \frac{f(M) - f(m)}{M - m}(B + C) + 2 \frac{f(m)M - f(M)m}{M - m} \\ &\leq \frac{f(M) - f(m)}{M - m}(A + D) + 2 \frac{f(m)M - f(M)m}{M - m} \quad (\text{by (i)}) \\ &= \frac{f(M) - f(m)}{M - m}A + \frac{f(m)M - f(M)m}{M - m} \\ &\quad + \frac{f(M) - f(m)}{M - m}D + \frac{f(m)M - f(M)m}{M - m} \\ &\leq f(A) + f(D) \quad (\text{by (2.4) and (2.5)}) \end{aligned}$$

The other cases are verified similarly. \square

As an immediate consequence of Theorem 2.10, we have

Corollary 2.11. *Let f and A, B, C, D be as in Theorem 2.10. If f is convex and one of the following conditions*

- (i) $B + C \leq A + D$ and $f(m) \leq f(M)$
- (ii) $A + D \leq B + C$ and $f(M) \leq f(m)$

is satisfied, then

$$f(B) + f(C) \leq g(A) + g(D),$$

for every continuous function $g \geq f$ on J and

$$g(B) + g(C) \leq f(A) + f(D),$$

or every continuous function $g \leq f$ on J .

Applying the above corollary to the power functions we get

Corollary 2.12. *Let $A, B, C, D \in \mathbb{B}(\mathcal{H})_h$ such that $I \leq A \leq m \leq B, C \leq M \leq D$ for two real numbers $m \leq M$. If one of the following conditions*

- (i) $B + C \leq A + D$ and $p \geq 1$
- (ii) $A + D \leq B + C$ and $p \leq 0$

is satisfied, then

$$B^p + C^p \leq A^q + D^q$$

for each $q \geq p$.

REFERENCES

1. T. Ando and X. Zhan, *Norm inequalities related to operator monotone functions*, Math. Ann. **315** (1999), 771–780.
2. K.M.R. Audenaert and J.S. Aujla, *On norm sub-additivity and super-additivity inequalities for concave and convex functions*, Linear Multilinear Algebra, DOI 10.1080/03081087.2011.653642.
3. J.S. Aujla and J.-C. Bourin, *Eigenvalue inequalities for convex and log-convex functions*, Linear Algebr and Appl. **424**(2007) 25–35.
4. J.S. Aujla and F. Silva, *Weak majorization inequalities and convex functions*, Linear Algebra Appl. **369** (2003), 217–233.
5. J.-C. Bourin and M. Uchiyama, *A matrix subadditivity inequality for $f(A+B)$ and $f(A)+f(B)$* , Linear Alg. Appl. **423** (2007) 512–518.
6. T. Furuta, J. Mićić Hot, J. Pečarić and Y. Seo, *Mond–Pečarić Method in Operator Inequalities*, Zagreb, Element, 2005.
7. T. Kosem, *Inequalities between $\|f(A+B)\|$ and $\|f(A)+f(B)\|$* , Linear Algebra Appl. **418** (2006), 153–160.
8. A. Matković, J. Pečarić and I. Perić, *A variant of Jensen’s inequality of Mercer’s type for operators with applications*, Linear Algebra Appl. **418** (2006), 551–564.
9. M.S. Moslehian and H. Najafi, *Around operator monotone functions*, Integral Equations Operator Theory **71** (2011), 575–582.
10. J. Pečarić, F. Proschan and Y.L. Tong, *Convex functions, partial orderings, and statistical applications*, Academic Press, Inc., Boston, MA, 1992.
11. O. Tychonov, *Subadditivity inequalities in von Neumann algebras and characterization of tracial functionals*, Positivity **9** (2005) 259–264.
12. M. Uchiyama, *Subadditivity of eigenvalue sums*, Proc. Amer. Math. Soc. **134** (2006), 1405–1412.

MOHAMMAD SAL MOSLEHIAN, MOHSEN KIAN: DEPARTMENT OF PURE MATHEMATICS, CENTER OF EXCELLENCE IN ANALYSIS ON ALGEBRAIC STRUCTURES (CEAAS), FERDOWSI UNIVERSITY OF MASHHAD, P.O. BOX 1159, MASHHAD 91775, IRAN.

E-mail address: moslehian@ferdowsi.um.ac.ir and moslehian@member.ams.org

E-mail address: kian.tak@yahoo.com

JADRANKA MIĆIĆ: FACULTY OF MECHANICAL ENGINEERING AND NAVAL ARCHITECTURE, UNIVERSITY OF ZAGREB, IVANA LUČIĆA 5, 10000 ZAGREB, CROATIA

E-mail address: jmicic@fsb.hr